

On co-edge-regular graphs with 4 distinct eigenvalues

Hong-Jun Ge^a and Jack H. Koolen^{*a,b}

^aSchool of Mathematical Sciences, University of Science and Technology of China, 96 Jinzhai Road, Hefei, 230026, Anhui, PR China

^bCAS Wu Wen-Tsun Key Laboratory of Mathematics, University of Science and Technology of China, 96 Jinzhai Road, Hefei, Anhui, 230026, PR China

June 29, 2026

Abstract

Tan et al. conjectured that connected co-edge-regular graphs with four distinct eigenvalues and fixed smallest eigenvalue, when having sufficiently large valency, belong to two different families of graphs. In this paper we construct two new infinite families of connected co-edge-regular graphs with four distinct eigenvalues and fixed smallest eigenvalue, thereby disproving their conjecture. Moreover, one of these constructions demonstrates that clique-extensions of Latin Square graphs are not determined by their spectrum.

1 Introduction

For undefined notions we refer to the next section and [3, 4, 5].

Over the past fifty years, strongly regular graphs have attracted a lot of attention, see [4]. Notice that a connected non-complete regular graph G is strongly regular if and only if it has exactly three distinct eigenvalues. In particular, Neumaier classified strongly regular graphs with fixed smallest eigenvalue in [13].

Theorem 1.1 (cf. [13, Theorem 5.1]). *Let $\lambda \geq 2$ be an integer. Except for finitely many exceptions, any strongly regular graph with smallest eigenvalue $-\lambda$ is a Steiner graph, a Latin Square graph, or a complete multipartite graph.*

In this paper, we study co-edge-regular graphs with four distinct eigenvalues and fixed smallest eigenvalue. We will show that for this class of graphs a result like Theorem 1.1 is difficult to obtain.

Notice that clique-extensions of strongly regular graphs are co-edge-regular graphs with at most four distinct eigenvalues. Hayat et al. [9] and Tan et al. [15] gave spectral characterizations of clique-extensions of Square grid graphs and triangular graphs under the assumption that graphs are co-edge-regular.

Theorem 1.2 (cf. [9, Theorem 1.1]). *Let G be a co-edge-regular graph with spectrum*

$$\left\{ (s(2t+1)-1)^1, (st-1)^{2t}, (-1)^{(s-1)(t+1)^2}, (-s-1)^{t^2} \right\},$$

where $s \geq 2$ and $t \geq 1$ are integers. If $t \geq 11(s+1)^3(s+2)$, then G is the s -clique extension of the $(t+1) \times (t+1)$ -grid.

*Corresponding author

2020 Mathematics Subject Classification. Primary: 05C50; Secondary: 05B15.

E-mail addresses: ghj17000225@mail.ustc.edu.cn (H.-J. Ge), koolen@ustc.edu.cn (J.H. Koolen).

Theorem 1.3 (cf. [15, Theorem 1]). *Let G be a co-edge-regular graph with spectrum*

$$\left\{ (2st - 3s - 1)^1, (st - 3s - 1)^{t-1}, (-1)^{\frac{(s-1)t(t-1)}{2}}, (-s - 1)^{\frac{t(t-3)}{2}} \right\},$$

where $s \geq 2$ and $t \geq 1$ are integers. If $t \geq 48s$, then G is the s -clique extension of the triangular graph $T(t)$.

Theorems 1.2 and Theorem 1.3 suggest that similar results may hold in the general case. Based on these results Tan et al. [15] conjectured the following:

Conjecture 1.4 (cf. [15, Conjecture 3]). *Let G be a connected k -regular graph with n vertices and co-edge-regular with parameters μ having four distinct eigenvalues. Let $t \geq 2$ be an integer. There exists a constant n_t such that, if $\theta_{\min}(G) \geq -t$, $n \geq n_t$ and $k < n - 2 - \frac{(t-1)^2}{4}$, then either G is the s -clique extension of a strongly regular graph for $2 \leq s \leq t - 1$ or G is a $p \times q$ -grid with $p > q \geq 2$.*

However, we will show that Conjecture 1.4 is false, by providing two infinite families of co-edge-regular graphs. This demonstrates that the class of co-edge-regular graphs is much more complicated than previously thought. The first family is based on results of Haemers and Tonchev [8]. The graphs in this family do not have -1 as an eigenvalue.

The second family is much more surprising since they have -1 as an eigenvalue and cospectral with certain clique-extensions of certain Latin Square graphs. In order to state the following result we will need to introduce co-edge-regular graphs of level t . For a pair x, y of adjacent vertices in a graph G , let $\lambda(x, y)$ be the number of common neighbours of x and y . We say that a co-edge-regular graph is of *level t* if $\#\{\lambda(x, y) \mid x, y \text{ are adjacent vertices}\} = t$. Note that, in the literature, see for example [6], the notion of a quasi-strongly regular graph of grade t exists. Such a graph is just the complement of a co-edge-regular graph of level t .

Theorem 1.5. *Let q be a prime power. For any positive integer $s \geq 2$, such that q is a factor of s , there are infinitely many integers n for which there exists a co-edge-regular graph with level at least 4 exists that is cospectral with the s -clique extension of each Latin Square graph $LS_{q+1}(qn)$ and hence has exactly four distinct eigenvalues.*

Note that the s -clique extension of any non-complete connected strongly regular graph has exactly four distinct eigenvalues and is co-edge-regular with level 2.

Theorem 1.5 demonstrates that to give a spectral characterization of the clique-extensions of strongly regular graphs with smallest eigenvalue at most -3 is impossible, even if one assumes that the graphs are co-edge-regular.

This paper is organized as follows. In Section 2, we provide definitions and preliminaries. Section 3 presents a combinatorial characterization of connected co-edge-regular graphs with exactly four distinct eigenvalues. In Section 4, we discuss a result from Haemers and Tonchev [8] and, as a consequence, construct an infinite family of connected co-edge-regular graphs of level 2 with exactly four distinct eigenvalues. Finally, in Section 5, we will construct co-edge-regular graphs of level 3 that are cospectral certain clique-extensions of certain Latin Square graphs. The existence of this family shows Theorem 1.5.

2 Definitions and Preliminaries

2.1 Graphs

A graph G is an ordered pair $(V(G), E(G))$, where $V(G)$ is a finite set and $E(G) \subseteq \binom{V(G)}{2}$. The set $V(G)$ (resp. $E(G)$) is called the vertex set (resp. edge set) of G , and the cardinality of $V(G)$ (resp. $E(G)$) is called the order (resp. size) of G and is denoted by $n(G)$ (resp. $e(G)$). The adjacency matrix of G , denoted by $A(G)$, is a symmetric $(0, 1)$ -matrix indexed by $V(G)$, such that

$(A(G))_{xy} = 1$ if and only if xy is an edge in G . The eigenvalues of G are the eigenvalues of $A(G)$, and the spectral radius, i.e. the largest eigenvalue, of G is denoted by $\rho(G)$. Two graphs are called *cospectral* if they have the same spectrum.

If x is a vertex of G then $N_G(x) := \{y \in V(G) \mid x \sim y\}$. The subgraph induced on $N_G(x)$ denoted by $\Delta_G(x)$, is called the local graph at x . If it is clear what the graph G is, we omit G in the notation. A *clique* (resp. *co-clique*) in a graph G is a set of vertices such that each pair of distinct vertices in it are adjacent (resp. non-adjacent). We sometimes consider a clique (resp. a co-clique) of G also as an induced subgraph of G .

Let G be a graph. Let $\pi := \{V_1, \dots, V_m\}$ be a partition of vertex set $V(G)$, and, for $1 \leq i, j \leq m$ and $u \in V_i$, let $b_{ij}(u)$ be the number of neighbors of u in V_j . Then partition π is called *equitable* if $b_{ij}(u)$ is independent from the concrete choice of $u \in V_i$. In this case, the $m \times m$ matrix $B := (b_{ij})$ is called the *quotient matrix* of π . It is well known that for an equitable partition of G with quotient matrix B , the eigenvalues of B are also eigenvalues of $A(G)$, see for example [5, Theorem 9.3.3].

2.2 Edge-regular and co-edge-regular graphs

For an edge xy in a graph G we denote by $\lambda(x, y)$, the number of common neighbours of x and y in G . For two non-adjacent vertices x and y we denote by $\mu(x, y)$, the number of common neighbours of x and y in G .

A regular graph is *edge-regular* with parameter λ if $\lambda(x, y) = \lambda$ for all edges xy of G . An edge-regular graph has *level* t if $\#\{\mu(x, y) \mid x, y \text{ are non-adjacent distinct vertices}\} = t$. This notion is also known as a quasi-strongly regular graph with grade t , see for example [6].

A regular graph is *co-edge-regular* with parameter μ if any two distinct non-adjacent vertices have exactly μ common neighbors. We say that a co-edge-regular graph has *level* t if $\#\{\lambda(x, y) \mid x, y \text{ adjacent vertices}\} = t$. Note that the complement of an edge-regular graph is co-edge-regular and vice versa and that the complement of an edge-regular graph which has level t is co-edge-regular and has level t and vice versa.

A graph G is called *strongly regular* with parameters (n, k, λ, μ) if it has n vertices, is k -regular, is edge-regular with parameter λ and co-edge-regular with parameter μ .

The following result is well-known, see [5, Lemma 10.2.1].

Lemma 2.1. *A connected non-complete regular graph G is strongly regular if and only if it has exactly three distinct eigenvalues.*

This means that a connected non-complete regular graph is a co-edge-regular graph of level 1 if and only if it has exactly three distinct eigenvalues.

Goldberg [6] conjectured a possible extension of a result from Jurišić et al. [11] who showed it for distance-regular graphs.

Conjecture 2.2 (cf. [6, Conjecture 1]). *Let G be a k -regular and edge-regular graph with parameter λ , and let θ, θ' be two distinct eigenvalues of G , different from k . Then*

$$\left(\theta + \frac{k}{\lambda + 1}\right)\left(\theta' + \frac{k}{\lambda + 1}\right) \geq -\frac{k\lambda(k - \lambda - 1)}{(\lambda + 1)^2}.$$

However, it is worth mentioning that, for any integer $s \geq 2$ and any integer $t \geq 3$, the complement of the s -clique extension of a Latin Square graph $LS_t(n)$ (see Subsection 2.5 for a definition) disproves this conjecture, if n is sufficiently large. Moreover, the complement of our construction in Section 5 also provides counterexamples to this conjecture.

2.3 Designs

A pair $(\mathcal{P}, \mathcal{B})$ is a 2 -(v, k, λ)-*design* if the following conditions are satisfied:

- (i) $|\mathcal{P}| = v$;

- (ii) For all $B \in \mathcal{B}$, we have $B \subseteq \mathcal{P}$ and $|B| = k$;
- (iii) For each pair of distinct elements x, y of $\mathcal{P}$ there are exactly λ elements B of $\mathcal{B}$ such that both x and y are in B .

We call the elements of $\mathcal{P}$ *points* and the elements of $\mathcal{B}$ *blocks*.

A 2 -(v, k, λ)-*design* $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ is called *resolvable* if we can partition the block set $\mathcal{B}$ into parts $\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_t$ such that $\mathcal{B}_i$ partition $\mathcal{P}$ for all $i = 1, 2, \dots, t$. We call the partition $\{\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_t\}$ a *resolution* of $\mathcal{D}$.

The *block graph* of a 2 -($v, k, 1$)-design $(\mathcal{P}, \mathcal{B})$ is the graph with vertex set $\mathcal{B}$ and two distinct blocks are adjacent if they intersect. Note that the block graph of a 2 -($v, k, 1$)-design is a strongly regular graph with smallest eigenvalue $-k$, see [4, Section 8.5.4A].

2.4 Association schemes

In this section, we introduce the definition of symmetric association schemes. We will not introduce the Bose-Mesner algebra for them. The reader is referred to [1, 5] for more information.

Let X be a finite set and let $\{R_0, R_1, \dots, R_D\}$ be a set of non-empty symmetric binary relations on X which partition $X \times X$. For any i ($0 \leq i \leq D$), define the matrix A_i with entries 0 and 1 such that $(A_i)_{xy} = 1$ if and only if $(x, y) \in R_i$. The pair $\mathfrak{X} = (X, \{R_i\}_{i=0}^D)$ is called a (*symmetric*) *association scheme with D classes* if the following conditions hold:

- (i) $A_0 = I_{|X|}$, which is the identity matrix of order $|X|$,
- (ii) $\sum_{i=0}^D A_i = J_{|X|}$, the all-ones matrix of order $|X|$,
- (iii) $A_i^\top = A_i$ for all $i \in \{0, 1, \dots, D\}$, where $A_i^\top$ is the transpose of A_i ,
- (iv) $A_i A_j = \sum_{h=0}^D p_{ij}^h A_h$, where p_{ij}^h are nonnegative integers, such that for all $x, y \in X$ with $(x, y) \in R_h$ the number of $z \in X$ with $(x, z) \in R_i$ and $(z, y) \in R_j$ equals p_{ij}^h .

We call R_0 the *trivial* relation and the other relations are called *non-trivial*. The integers p_{ij}^h are called the *intersection numbers* of $\mathfrak{X}$. Any association scheme in this paper is a symmetric association scheme. We also call $\mathfrak{X}$ a *D -class association scheme*. The matrices A_i , $i = 0, 1, \dots, D$ are called the *relation matrices* of the association scheme $\mathfrak{X}$, and the graph $G_i = (X, R_i)$, with adjacency matrix A_i , is called the *relation graph* for the relation R_i for $i = 0, 1, \dots, D$. Note that G_i is k_i -regular where $k_i = p_{ii}^0$, and we say that k_i is the valency of the relation R_i . Note that any relation graph in an association scheme $\mathfrak{X}$ with D classes has at most $D + 1$ distinct eigenvalues.

Van Dam gave a characterization of edge-regular graphs of level 2 with four distinct eigenvalues in [18].

Theorem 2.3 (cf. [18, Theorem 5.1]). *Let G be a connected regular graph that has four distinct eigenvalues. Then G is edge-regular of level 2 if and only if G is one of the classes of a 3-class symmetric association scheme.*

2.5 Orthogonal arrays

An *orthogonal array* O of order (n, t) , denoted by $OA(n, t)$, is a $t \times n^2$ array such that each entry is an element of $[n]$ and $\#\{(r_i(\ell), r_j(\ell)) \mid \ell = 1, 2, \dots, n^2\} = n^2$ for $1 \leq i < j \leq t$, where r_i is the i -th row of O . Note that from an $OA(n, t)$, we can construct $t - 2$ mutually orthogonal Latin Squares of order n for $t \geq 3$, and vice versa. A *group-divisible orthogonal array* O of order (n, s, t) , denoted by $GOA(n, s, t)$ with groups $G_1, G_2, \dots, G_t$, each consisting of distinct s rows is an $st \times n^2$ -array such that $\#\{(r_i(\ell), r_j(\ell)) \mid \ell = 1, 2, \dots, n^2\} = n^2$, whenever r_i belongs to a different group than r_j . We allow repeated rows in a group.

If there exists an $OA(n, t)$, then, for all $s \geq 2$, there exists a $GOA(n, s, t)$.

The following theorem is the well known MacNeish's Theorem [12].

Theorem 2.4. If $n = p_1^{\alpha_1} \cdots p_r^{\alpha_r}$ where $p_1, \dots, p_r$ are distinct primes. Then there exists an $OA(n, r)$ with $r = \min_i p_i^{\alpha_i} + 1$

Now we will define the Latin Square graphs.

Definition 2.5. Let $\mathcal{O}$ be an $OA(n, m)$ with columns $c_1, c_2, \dots, c_{n^2}$. The Latin Square graph $LS_m(n)$ with respect to $\mathcal{O}$ is the graph on $V = \{c_1, c_2, \dots, c_{n^2}\}$ and for $i \neq j$, $c_i \sim c_j$ if there exists an integer ℓ such that $c_i(\ell) = c_j(\ell)$.

Each Latin Square graph $LS_m(n)$ is a strongly regular graph with parameters

$$(n^2, (n-1)m, (m-1)(m-2) + n - 2, m(m-1))$$

and has spectrum $\text{Spec}(LS_m(n)) = \{[(n-1)m]^1, [n-m]^{(n-1)m}, [-m]^{(n-1)(n+1-m)}\}$. For more background about Latin Square graphs, we refer to [4, 5]

For a positive integer s , the s -clique extension of a graph G is the graph $\tilde{G}$ obtained from G by replacing each vertex $x \in V(G)$ by a clique $\tilde{X}$ with s vertices, such that $\tilde{x} \sim \tilde{y}$ (for $\tilde{x} \in \tilde{X}$, $\tilde{y} \in \tilde{Y}$) in $\tilde{G}$ if and only if $x \sim y$ in G . Note that $\tilde{G}$ has adjacent matrix $A(\tilde{G}) = J_s \otimes (A(G) + I_v) - I_{sv}$, where I is the identity matrix, J is the all-ones matrix, and v is the number of vertices in G . In particular, if $\text{Spec}(G) = \{[\theta_0]^{m_0}, [\theta_1]^{m_1}, \dots, [\theta_r]^{m_r}\}$, then

$$\text{Spec}(\tilde{G}) = \{[s(\theta_0 + 1) - 1]^{m_0}, [s(\theta_1 + 1) - 1]^{m_1}, \dots, [s(\theta_r + 1) - 1]^{m_r}, [-1]^{(s-1)v}\}.$$

Note that if G and H have the same spectrum, then the s -clique extension of G has the same spectrum, as the s -clique extension of H . For more background about s -clique extensions, we refer to [9].

2.6 Parallel classes

Let q be a prime power and $\mathbb{F}_q$ be the finite field of order q . A plane P is an affine plane in $\mathbb{F}_q^3$. Two planes P_1 and P_2 are *parallel* if they are disjoint. A *parallel class* $\mathcal{S}$ is a set of q mutually parallel planes in $\mathbb{F}_q^3$. Bose first considered these sets in [2] to construct certain designs. In the next result we show that there are special parallel classes in $\mathbb{F}_q^3$. The following result may be known, but we could not find a reference for it.

Theorem 2.6. There are $q + 1$ parallel classes $\mathcal{S}_1, \dots, \mathcal{S}_{q+1}$ in $\mathbb{F}_q^3$ satisfying:

- (i) $|P \cap Q| = q$ for $P \in \mathcal{S}_i$ and $Q \in \mathcal{S}_j$ if $i \neq j$,
- (ii) $|P \cap Q \cap R| = 1$ for $P \in \mathcal{S}_h$, $Q \in \mathcal{S}_i$ and $R \in \mathcal{S}_j$, if $1 \leq h < i < j \leq q + 1$.

Proof. Let $v_\infty := (0, 0, 1)$ and $v_x := (1, x, x(x+1))$ for each $x \in \mathbb{F}_q$. Let x, y, z be three distinct elements of $\mathbb{F}_q$. Then

$$\det \begin{pmatrix} 0 & 0 & 1 \\ 1 & x & x(x+1) \\ 1 & y & y(y+1) \end{pmatrix} = y - x$$

and

$$\det \begin{pmatrix} 1 & x & x(x+1) \\ 1 & y & y(y+1) \\ 1 & z & z(z+1) \end{pmatrix} = (x-y)(y-z)(z-x).$$

This means that any three distinct vectors of $\{v_x \mid x \in \mathbb{F}_q\} \cup \{v_\infty\}$ are linear independent. Let $P_a^b := \{u \in \mathbb{F}_q^3 \mid u^T v_a = b\}$ for $a \in \mathbb{F}_q \cup \{q+1\}$ and $b \in \mathbb{F}_q$. Then $\mathcal{S}_a := \{P_a^b \mid b \in \mathbb{F}_q\}$ is a parallel class as these planes are all orthogonal to v_a . Let $a_1, a_2, a_3 \in \mathbb{F}_q \cup \{v_\infty\}$ be pairwise distinct. Then we have $|P_{a_1}^{b_1} \cap P_{a_2}^{b_2}| = q$ for $b_1, b_2 \in \mathbb{F}_q$, as v_{a_1} and v_{a_2} are linear independent, and $|P_{a_1}^{b_1} \cap P_{a_2}^{b_2} \cap P_{a_3}^{b_3}| = 1$ for $b_1, b_2, b_3 \in \mathbb{F}_q$, as v_{a_1}, v_{a_2} and v_{a_3} are linear independent. □

3 A combinatorial characterization of co-edge-regular graphs with four distinct eigenvalues

In this section we will study connected co-edge-regular graphs with at most four distinct eigenvalues. Note that by Lemma 2.1, we have a characterization of connected regular graphs with 3 distinct eigenvalues. We will generalize this to four distinct eigenvalues.

Let G be a connected k -regular graph on n vertices with exactly four eigenvalues $\theta_0 = k > \theta_1 > \theta_2 > \theta_3$. Let $A := A(G)$ be the adjacent matrix of G , $n := |V(G)|$, and $\ell := \frac{(k-\theta_1)(k-\theta_2)(k-\theta_3)}{n}$. Hoffman [10], see also [17], showed $(A - \theta_1 I)(A - \theta_2 I)(A - \theta_3 I) = \ell J$. It follows that

$$A^3 - (\theta_1 + \theta_2 + \theta_3)A^2 + (\theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3)A - \theta_1\theta_2\theta_3 I = \ell J. \quad (1)$$

Definition 3.1. (i) A co-edge-regular graph with parameter μ is a (μ, γ) -strongly co-edge-regular, if every pair of non-adjacent vertices $\{x, y\}$ satisfies $\sum_{z \in N(x) \cap N(y)} \lambda(x, z) = \gamma$.

(ii) A regular graph is (α, β) -weakly edge-regular if every pair of adjacent vertices $\{x, y\}$ satisfies $\alpha\lambda(x, y) = \sum_{z \in N(x) \cap N(y)} \lambda(x, z) + \beta$.

Note that a non-complete strongly regular graph with parameters (n, k, λ, μ) is (μ, γ) -strongly co-edge-regular with $\gamma = \lambda^2$, and (α, β) -weakly edge-regular for any pair (α, β) satisfying $\beta = (\alpha - \lambda)\lambda$.

Theorem 3.2. Let G be a connected non-complete k -regular and co-edge-regular graph on n vertices with parameter μ . Then G has at most 4 distinct eigenvalues if and only if it is (α, β) -weakly edge-regular and (μ, γ) -strongly co-edge-regular for some reals α, β, γ . Furthermore, if G has exactly 4 distinct eigenvalues $k > \theta_1 > \theta_2 > \theta_3$, then $\alpha - \mu = \theta_1 + \theta_2 + \theta_3$, $\mu(\alpha - 1) + k - \beta - \gamma = -(\theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3)$, and $\mu(k - \alpha) + \gamma = \frac{(k-\theta_1)(k-\theta_2)(k-\theta_3)}{n}$.

Proof. Let G be a k -regular and co-edge-regular graph with parameter μ .

As G is non-complete, G has at least three distinct eigenvalues. If G has exactly three distinct eigenvalue, then we have already seen that G is strongly and hence (α, β) -weakly edge-regular and (μ, γ) -strongly co-edge-regular for some reals α, β, γ . Now assume that G has at least four distinct eigenvalues.

Notice that $(A^3)_{xx} = 2e(\Delta_x) = k(k-1) - \mu(n-k-1)$, where $e(\Delta_x)$ is the number of edges in Δ_x the local graph at x . For a vertex x of G , let $N(x)$ be the set of neighbours of x in G .

Let x and y be two distinct non-adjacent vertices. Then

$$(A^3)_{xy} = (k - \mu)\mu + \sum_{z \in N(x) \cap N(y)} \lambda(x, z) \text{ and } (A^2)_{xy} = \mu.$$

Similarly, if u and v are two adjacent vertices, then

$$(A^3)_{uv} = \sum_{w \in N(u) \cap N(v)} \lambda(u, w) + (k - 1 - \lambda(u, v))\mu + k \text{ and } (A^2)_{uv} = \lambda(u, v).$$

Assume that G has exactly four distinct eigenvalues $\theta_0 = k > \theta_1 > \theta_2 > \theta_3$. By Equation (1), for two distinct non-adjacent vertices x and y , the sum

$$\sum_{z \in N(x) \cap N(y)} \lambda(x, z) = \mu(\theta_1 + \theta_2 + \theta_3 - k + \mu) + \ell$$

does not depend on the choice of x and y , where $\ell := \frac{(k-\theta_1)(k-\theta_2)(k-\theta_3)}{n}$. Hence, G is strongly co-edge-regular with $\gamma = \mu(\theta_1 + \theta_2 + \theta_3 - k + \mu) + \ell$.

Also by Equation (1), for two adjacent vertices u and v , the sum

$$(\theta_1 + \theta_2 + \theta_3 + \mu)\lambda(u, v) = \sum_{w \in N(u) \cap N(v)} \lambda(u, w) + (k - 1)\mu + k + \theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3 - \ell$$

does not depend on the choice of u and v . Hence, G is weakly edge-regular with $\alpha = \theta_1 + \theta_2 + \theta_3 + \mu$ and $\beta = (k-1)\mu + k + \theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3 - \ell$.

Now let G be a (μ, γ) -strongly co-edge-regular and (α, β) -weakly edge-regular graph. Let $f_1 := \mu(\alpha - 1) + k - \beta - \gamma$, $f_2 := \alpha - \mu$, $f_3 := k(k-1-\alpha) + \mu\alpha - \gamma - \mu(n-k-1)$, and $f_4 := \mu(k-\alpha) + \gamma$.

Let x be a vertex of G . Then $(A^3)_{xx} = k(k-1) - \mu(n-k-1) = f_3 + kf_2 + f_4 = (f_1A + f_2A^2 + f_3I + f_4J)_{xx}$.

Let x and y be two distinct non-adjacent vertices. Then

$$(A^3)_{xy} = (k-\mu)\mu + \sum_{z \in N(x) \cap N(y)} \lambda(x, z) \text{ and } (A^2)_{xy} = \mu.$$

As G is a (μ, γ) -strongly co-edge-regular graph, we see that $(A^3)_{xy} = (k-\mu)\mu + \gamma = f_4 + f_2\mu = f_4J_{xy} + f_2(A^2)_{xy} = (f_1A + f_2A^2 + f_3I + f_4J)_{xy}$.

Using the fact that G is (α, β) -weakly edge-regular, it can be checked in a similar way that $(A^3)_{uv} = (f_1A + f_2A^2 + f_3I + f_4J)_{uv}$ for adjacent vertices u, v of G . This means that $A^3 = f_1A + f_2A^2 + f_3I + f_4J$. This implies that $A^4 - (k+f_1)A^3 + (kf_1-f_2)A^2 + (kf_2-f_3)A + kf_3I = 0$, since $AJ = kJ$. Therefore, the degree of the minimal polynomial of A is at most four, and thus G has at most 4 distinct eigenvalues. $\square$

Of course the above theorem also gives a characterization of edge-regular graphs with four distinct eigenvalues, by looking at their complements. Note that this characterization generalizes the following result of Van Dam [18], as relation graphs from d -class association schemes have at most $d+1$ distinct eigenvalues.

Theorem 3.3 (cf. [18, Theorem 5.1]). *Let G be an edge-regular graph of level 2. Then G is the relation graph of a 3-class association scheme if and only if it has exactly four distinct eigenvalues.*

4 On a theme of Haemers and Tonchev

In strongly regular graphs, the cliques and co-cliques with the property that every vertex outside is adjacent with the same number of vertices inside are characterized by the fact that they satisfy with the equality so-called *Hoffman bound*:

Theorem 4.1 (cf. [4, Proposition 1.1.7]). *Let G be a strongly regular graph with parameters (n, k, λ, μ) and smallest eigenvalue $-m$.*

- (i) *If C is a clique of G , then $|C| \leq \frac{m+k}{m}$, with equality if and only if every vertex $x \notin C$ has the same number $\frac{\mu}{m}$ of neighbors in C .*
- (ii) *If D is a co-clique of G , then $|D| \leq \frac{m}{m+k}n$, with equality if and only if every vertex $x \notin D$ has the same number m of neighbors in D .*
- (iii) *If a clique C and a co-clique D both meet the bounds of (i) and (ii), then $|C \cap D| = 1$.*

In 2021, Haemers wrote a note [7] clarifying the history of the Hoffman bound. We call a (co-)clique that meets the Hoffman bound a *Hoffman (co-)clique*. Note that in the literature a Hoffman clique is also called a *Delsarte clique*.

Definition 4.2. (i) *A Hoffman coloring in a strongly regular graph G is a partition of the vertex set into Hoffman co-cliques;*

- (ii) *A Hoffman spread in a strongly regular graph G is a partition of the vertex set into Hoffman cliques.*

Note that a Hoffman coloring in a strongly regular graph G is the same as a Hoffman spread in its complement $\overline{G}$.

In [8], Haemers and Tonchev showed the following result, generalizing a result of Brouwer.

Theorem 4.3 (cf. [8, Proposition 4.1]). *Let G be a connected non-complete strongly regular graph with n vertices and spectrum $\text{Spec}(G) = \{[k]^1, [r]^f, [s]^g\}$ having a Hoffman spread $S = \{C_1, C_2, \dots, C_t\}$. Let E be the edge set of G and F_i be the edge set of C_i for $i = 1, 2, \dots, t$. Let G' be the graph with the same vertex set as G and edge set $E \setminus \bigcup_{i=1}^t F_i$. Then G' is the relation graph of a 3-class association scheme. In particular, G' is regular and edge-regular with spectrum*

$$\text{Spec}(G') = \{[k + \frac{n-k-1}{r+1}]^1, [r-1]^{m_1}, [s-1]^{m_2}, [s + \frac{n-k-1}{r+1}]^{m_3}\},$$

where $m_1 = f - \frac{sn}{s-k} + 1$, $m_2 = g$, and $m_3 = \frac{sn}{s-k} - 1$.

Note that the complement $\overline{G_1}$ of G_1 of Theorem 4.3, is co-edge-regular and has at most four distinct eigenvalues. Therefore, each strongly regular graph with a Hoffman spread can construct a counterexample of Conjecture 1.4. There are many examples of strongly regular graphs with Hoffman spreads, see [8, 16].

By using resolvable $2-(v, t, 1)$ -designs, we give an infinite family of counterexamples of Conjecture 1.4 by the above theorem. The complements of these examples were obtained by Haemers and Tonchev [8]. For the convenience of the reader, we give the details.

Let $\mathcal{D} = (\mathcal{P}, \mathcal{B})$ be a resolvable $2-(v, t, 1)$ -design, say, with resolution $\rho = \{\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_\ell\}$. This means that the block graph G of $\mathcal{D}$ is a strongly regular graph with parameters (n, k, λ, μ) having smallest eigenvalue $-t$, see [4, Section 8.5.4A], where $n = \frac{v(v-1)}{t(t-1)}$, $k = \frac{t(v-t)}{t-1}$, $\lambda = t(t-2) + \frac{v-t}{t-1}$, and $\mu = t^2$. Note that each $\mathcal{B}_i$ is a co-clique of G with order $\frac{v}{t}$. By Theorem 4.1, each $\mathcal{B}_i$ denotes a Hoffman co-clique of G . Hence, the resolution ρ gives a Hoffman coloring of G .

Now construct the H with vertex set $\mathcal{B}$ such that, for two distinct blocks B_1, B_2 , we have $B_1 \sim B_2$ if they were adjacent in G or there exists i such that $B_1, B_2 \in \mathcal{B}_i$. Then, by Theorem 4.3, we see that H is a co-edge-regular graph with smallest eigenvalue $-t-1$ having exactly four distinct eigenvalues.

Theorem 4.4. *The graph H is regular with valency $\frac{v-t}{t-1}(t + \frac{t-1}{t})$ and co-edge-regular of level 2 with parameter $t(t+2)$. Moreover, the spectrum of H is*

$$\text{Spec}(H) = \{[\frac{v-t}{t-1}(t + \frac{t-1}{t})]^1, [\frac{v-t}{t} - t]^{f_1}, [\frac{v-1}{t-1} - t - 2]^{f_2}, [-t-1]^{f_3}\},$$

where $f_1 = \frac{v-t}{t-1}$, $f_2 = v-1$, and $f_3 = \frac{v-1}{t}(\frac{v-t}{t-1} - t)$.

Proof. By Theorem 2.3 and Theorem 4.3, it suffices to show that H is co-edge-regular with parameter $t(t+2)$. Let $x, y \in V(H)$ be two non-adjacent vertices. Without loss of generality, assume that $x \in \mathcal{B}_1$ and $y \in \mathcal{B}_2$. Since $\mathcal{B}_i$ is a Hoffman co-clique of G for each i , Theorem 4.1 implies that x (resp. y) has exactly t neighbors in $\mathcal{B}_2$ (resp. $\mathcal{B}_1$). Note that G is co-edge-regular with parameter t^2 . Therefore, $|N_H(x) \cap N_H(y)| = |N_G(x) \cap N_G(y)| + |N_H(x) \cap \mathcal{B}_2| + |N_H(y) \cap \mathcal{B}_1| = t(t+2)$. This completes the proof. $\square$

Note that for fixed $t \geq 2$, there are infinitely many v such that there exists a resolvable $2-(v, t, 1)$ -design, see for example [14]. These give infinitely many counterexamples to Conjecture 1.4.

5 Another construction

In this section, we construct a new family of co-edge-regular graphs with four distinct eigenvalues of which one is -1 . We will use group-divisible orthogonal arrays for this construction. We

will call these new graphs the twisted Latin Square graphs. In the rest of this section, let q be a prime power and n be a positive integer such that there exists a $GOA(n, q, q+1)$, say $\mathcal{O}$ with groups $G_1, G_2, \dots, G_{q+1}$. Note that, for a given prime power q , there are infinitely many n such that exists a $GOA(n, q, q+1)$, by MacNeish's Theorem [12]. For $i \in [q+1]$, order the rows in G_i as r_j^i where $j \in [q]$, since there are exactly q rows in G_i .

By Theorem 2.6, there are $q+1$ parallel classes $\mathcal{S}_1, \dots, \mathcal{S}_{q+1}$ in $\mathbb{F}_q^3$ satisfying:

- (i) $|P \cap Q| = q$ for $P \in \mathcal{S}_i$ and $Q \in \mathcal{S}_j$ if $i \neq j$,
- (ii) $|P \cap Q \cap R| = 1$ for $P \in \mathcal{S}_h$, $Q \in \mathcal{S}_i$ and $R \in \mathcal{S}_j$, if $1 \leq h < i < j \leq q+1$.

Let $\mathcal{S}_i = \{P_1^i, P_2^i, \dots, P_q^i\}$ for $i \in [q+1]$. We consider the planes P_j^i as subsets of order q^2 of $\mathbb{F}_q^3$ for $i \in [q+1]$ and $j \in [q]$.

Now we are ready to define the twisted Latin Square graphs. As we will see later they have the same spectra as certain clique-extensions of certain Latin Square graphs.

The *twisted Latin Square graph* with parameters (q, n) with respect to $\mathcal{O}$, abbreviated with $TLS(q, n)$, is the graph with vertex set $\mathbb{F}_q^3 \times [n^2]$. Two distinct vertices $(\mathbf{x}, i)$ and $(\mathbf{y}, j)$ of $TLS(q, n)$ are adjacent if $i = j$ or if there exists a plane P_ℓ^p such that $\mathbf{x}$ and $\mathbf{y}$ both lie on and $r_\ell^p(i) = r_\ell^p(j)$.

For $s \in [q+1]$, $t \in [q]$ and $\ell \in [n]$ define the set $C(s, t, \ell)$ as the set consisting of the vertices $(\mathbf{x}, i)$ such that $\mathbf{x} \in P_t^s$ and $r_t^s(i) = \ell$. It is clear that the set $C(s, t, \ell)$ induces a clique in $TLS(q, n)$ of order $q^2 n$.

5.1 Structure of $TLS(q, n)$

We denote $F(i) := \{(\mathbf{x}, i) \mid \mathbf{x} \in \mathbb{F}_q^3\}$ for $i = 1, 2, \dots, n^2$ and $P_t^s(i) := \{(\mathbf{x}, i) \mid \mathbf{x} \in P_t^s\}$ for $s \in [q+1]$, $t \in [q]$, and $i \in [n^2]$.

Lemma 5.1. *Let $s_1, s_2 \in [q+1]$, $t_1, t_2 \in [q]$ and $\ell_1, \ell_2 \in [n]$. Assume $C(s_1, t_1, \ell_1) \neq C(s_2, t_2, \ell_2)$. If $s_1 \neq s_2$, then there exists an integer i such that $C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2) = P_{t_1}^{s_1}(i) \cap P_{t_2}^{s_2}(i) \subseteq F(i)$. and $|C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2)| = q$. If $s_1 = s_2$, then $|C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2)| = 0$.*

Proof. Assume $s_1 \neq s_2$. By Theorem 2.6, the planes $P_{t_1}^{s_1}$ and $P_{t_2}^{s_2}$ intersect in q vectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_q$. There exists exactly one position $i \in [n^2]$ such that $r_{t_1}^{s_1}(i) = \ell_1$ and $r_{t_2}^{s_2}(i) = \ell_2$, by the definition of $\mathcal{O}$. This shows that $C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2) = P_{t_1}^{s_1}(i) \cap P_{t_2}^{s_2}(i)$ and $|C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2)| = q$ in this case. Now assume $s_1 = s_2$. If $t_1 \neq t_2$, then the planes $P_{t_1}^{s_1}$ and $P_{t_2}^{s_2}$ are parallel and disjoint, and thus $|C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2)| = 0$. If $t_1 = t_2$, then $\ell_1 \neq \ell_2$ and again it is clear that $|C(s_1, t_1, \ell_1) \cap C(s_2, t_2, \ell_2)| = 0$. This shows the lemma. $\square$

Lemma 5.2. *Any twisted Latin Square graph $TLS(q, n)$ is regular with valency $k = q^3 - 1 + (q+1)(n-1)q^2$.*

Proof. Let $(\mathbf{x}, i)$ be a vertex of $TLS(q, n)$. Then it has $q^3 - 1$ neighbours in $F(i)$. The vector $\mathbf{x}$ lies in the unique plane, say $P_{t_s}^s$, in the parallel class $\mathcal{S}_s$ for $s \in [q+1]$. So $(\mathbf{x}, i)$ has exactly $q^2(n-1)$ neighbours in $C(s, t_s, r_{t_s}^s(i)) - F(i) = C(s, t_s, r_{t_s}^s(i)) - P_{t_s}^s(i)$. If $s_1 \neq s_2$, then, by Lemma 5.1, $C(s_1, t_{s_1}, \ell_1) \cap C(s_2, t_{s_2}, \ell_2) = P_{t_{s_1}}^{s_1}(i) \cap P_{t_{s_2}}^{s_2}(i) \subset F(i)$, as $(\mathbf{x}, i)$ is one of them. It follows that the valency of $(\mathbf{x}, i)$ is exactly $q^3 - 1 + (q+1)(n-1)q^2$. $\square$

Lemma 5.3. *Let $s \in [q+1]$, $t \in [q]$ and $\ell \in [n]$. Let $v = (\mathbf{x}, i)$ be a vertex not in $C(s, t, \ell)$. Assume that $\mathbf{x}$ lies in planes $P_{t_j}^j$ for $j \in [q+1]$. If $r_t^s(i) = \ell$, then $C(s, t, \ell) \cap N(v) = P_t^s(i)$ holds. If $r_t^s(i) \neq \ell$, then there exists pairwise distinct m_p with $p \in [q+1] - \{s\}$ such that*

$$C(s, t, \ell) \cap N(v) = \bigcup_{p=1, p \neq s}^{q+1} P_{t_p}^p(m_p) \cap P_t^s(m_p).$$

In particular, v has exactly q^2 neighbours in $C(s, t, \ell)$.

Proof. Assume that $v = (\mathbf{x}, i)$. The vector $\mathbf{x}$ lies in a unique plane, say $P_{t_j}^j$, in the parallel class $\mathcal{S}_j$ for $j \in [q+1]$. Hence, $(\mathbf{x}, i)$ lies in the clique $C(j, t_j, r_{t_j}^j(i))$ for $j \in [q+1]$.

If $r_t^s(i) = \ell$, then $P_t^s(i) = C(s, t, \ell) \cap F(i) \subset C(s, t, \ell) \cap N(v)$. So v has at least q^2 neighbours in $C(s, t, \ell)$, and $t_s \neq t$ as otherwise $v \in C(s, t, \ell)$. As $(\mathbf{x}, i)$ lies in the clique $C(j, t_j, r_{t_j}^j(i))$ for $j \in [q+1]$ and, by Lemma 5.1, $C(s, t, \ell) \cap C(j, t_j, r_{t_j}^j(i)) = P_t^s(i) \cap P_{t_j}^j \subset P_t^s(i)$ if $j \neq s$. This means that $|C(s, t, \ell) \cap N(v)| = |P_t^s(i)| = q^2$ holds.

If $r_t^s(i) \neq \ell$, then $C(s, t, \ell) \cap N(v) = C(s, t, \ell) \cap (\bigcup_{i \neq s} C(j, t_j, r_{t_j}^j(i)))$. By Lemma 5.1, there exists an integer m_j such that $C(s, t, \ell) \cap C(j, t_j, r_{t_j}^j(i)) = P_t^s(m_j) \cap P_{t_j}^j(m_j)$ if $j \neq s$. As $r_t^s(i) \neq \ell$ and $r_t^s(m_j) = \ell$, we have $i \neq m_j$ for all $j \neq s$.

Now we show that $m_j \neq m_{j'}$ for distinct $j, j' \in [q+1] - \{s\}$. In fact, we have $(r_{t_j}^j(m_j), r_{t_{j'}}^{j'}(m_j)) \neq (r_{t_j}^j(i), r_{t_{j'}}^{j'}(i))$, by the definition of $\mathcal{O}$, as $i \neq m_j$. Thus, $m_j \neq m_{j'}$ for distinct $j, j' \in [q+1] - \{s\}$.

This implies that $|C(s, t, \ell) \cap N(v)| = |C(s, t, \ell) \cap (\bigcup_{i \neq s} C(j, t_j, r_{t_j}^j(i)))| = |\bigcup_{i \neq s} (P_t^s(m_j) \cap P_{t_j}^j(m_j))| = \sum_{i \neq s} |P_t^s(m_j) \cap P_{t_j}^j(m_j)| = q^2$ by Theorem 2.6. $\square$

Lemma 5.4. *Each twisted Latin Square graph $TSL(q, n)$ has $q^2(n-1) - 1$ as an eigenvalue.*

Proof. Let V be the vertex set of $TSL(q, n)$. By Lemma 5.2, $TSL(q, n)$ is regular with valency $k = q^3 - 1 + (q+1)(n-1)q^2$. By Lemma 5.3, for $s \in [q+1]$, we have a equitable partition $\{C(s, t, \ell), V - C(s, t, \ell)\}$ with quotient matrix $\begin{pmatrix} nq^2 - 1 & k - nq^2 + 1 \\ q^2 & k - q^2 \end{pmatrix}$ having eigenvalues $q^2(n-1) - 1$ and k . Therefore, $q^2(n-1) - 1$ is an eigenvalue of $TSL(q, n)$. $\square$

Lemma 5.5. *Let $u = (\mathbf{x}, i)$ be a vertex in $TSL(q, n)$ where $\mathbf{x} \in \mathbb{F}_q^3$ and $i \in [n^2]$. If u has neighbours in $F(m)$ where $m \neq i$ and $m \in [n^2]$. Then there exists $s \in [q+1]$ and $t \in [q]$ such that $F(m) \cap N(u) = P_t^s(m)$.*

Proof. Let $v = (\mathbf{y}, m)$ be a neighbour of u in $F(m)$. This means that there exists $s \in [q+1], t \in [q]$ and $\ell \in [n]$ such that $u, v \in C(s, t, \ell)$, which implies $\ell = r_t^s(i) = r_t^s(m)$. It follows that $N(u) \cap F(m) \supseteq P_t^s(m)$. Let $s_1 \in [q+1], t_1 \in [q]$ such that $s_1 \neq s$. Then $r_{t_1}^{s_1}(i) \neq r_{t_1}^{s_1}(m)$ by the definition of $\mathcal{O}$. As a consequence, we find that $N(u) \cap F(m) = P_t^s(m)$. This shows the lemma. $\square$

5.2 Local graphs of $TSL(q, n)$

Let $u = (\mathbf{x}, m)$ be a vertex of the graph $TSL(q, n)$, where $\mathbf{x} \in \mathbb{F}_q^3$ and $m \in [n^2]$. The vector $\mathbf{x}$ lies in the unique plane, say $P_{t_s}^s$, in the parallel class $\mathcal{S}_s$ for $s \in [q+1]$. This means that $(\mathbf{x}, m)$ lies in the clique $C(s, t_s, r_{t_s}^s(m))$ for $s \in [q+1]$. We define the set $A_{ij} = A_{ij}(u)$ as follows: Let $A_{ij} := P_{t_i}^i(m) \cap P_{t_j}^j(m) - \{(\mathbf{x}, m)\}$ for distinct $i, j \in [q+1]$.

For $i \in [q+1]$, we define the sets $B_i = B_i(u)$, $C_i = C_i(u)$ and $R = R(u)$ by $B_i := P_{t_i}^i(m) - \bigcup_{j \neq i} P_{t_j}^j(m)$, $C_i := C(i, t_i, r_{t_i}^i(m)) - P_{t_i}^i(m)$, and $R := F(m) - \bigcup_{j=1}^{q+1} P_{t_j}^j(m)$.

We first give the cardinalities of these sets and also show that they are mutually disjoint.

Proposition 5.6. *Let A_{ij} , B_i , C_i , and R be the sets as defined above. These sets satisfy the following properties:*

- (i) *The sets A_{ij} , B_h , C_p , and R are mutually disjoint for $i, j, h, p \in [q+1]$ and $i \neq j$;*
- (ii) *$|A_{ij}| = q - 1$ for $i, j \in [q+1]$ and $i \neq j$;*
- (iii) *$|B_i| = q - 1$ for $i \in [q+1]$;*
- (iv) *$|C_i| = (n-1)q^2$ for $i \in [q+1]$;*
- (v) *$|R| = \frac{q(q-1)^2}{2}$ for $i \in [q+1]$.*

Proof. As A_{ij}, B_h, R are all subsets of $F(m)$ and $C_p \cap F(m) = \emptyset$ for $i, j, h, p \in [q+1]$, $i \neq j$ so C_p is disjoint from B_h, R and A_{ij} . For distinct $i, j \in [q+1]$, by Lemma 5.1, we have

$$C(i, t_i, r_{t_i}^i(m)) \cap C(j, t_j, r_{t_j}^j(m)) \subset F(m).$$

So $C_i \cap C_j = \emptyset$. Clearly the B_i and B_j are disjoint if $i \neq j$, and $A_{ij} \cap B_h = \emptyset$ if $i \neq j$. It is also clear that the set R is disjoint from the sets A_{ij} and B_h . Let $i_1, i_2, j_1, j_2 \in [q+1]$ such that $i_1 \neq j_1$, $i_2 \neq j_2$ and $\{i_1, j_1\} \neq \{i_2, j_2\}$. Then $A_{i_1, j_1} \cap A_{i_2, j_2}$ is contained in the intersection of at least three planes all containing $(\mathbf{x}, m)$. This means that $A_{i_1, j_1} \cap A_{i_2, j_2} = \emptyset$, by Theorem 2.6. These show that all the sets in (i) are disjoint.

Now we will determine the order of the sets A_{ij}, B_i, C_i and R . For $i \neq j \in [q+1]$ the set $A_{ij} = P_{t_i}^i(m) \cap P_{t_j}^j(m) \setminus u$, and hence $|A_{ij}| = q-1$ by Theorem 2.6. It follows that $|B_i| = |P_{t_i}^i| - \sum_{j \neq i} |A_{ij}| - 1 = q^2 - q(q-1) - 1 = q-1$ for $i \in [q+1]$, and thus

$$|R| = q^3 - \sum_i |B_i| - \sum_{i=1}^{q+1} \sum_{j=i+1}^{q+1} |A_{ij}| - |\{u\}| = q^3 - (q+1)(q-1) - \frac{(q+1)q}{2}(q-1) - 1 = \frac{q(q-1)^2}{2}.$$

Note that $|C_i| = |C(i, t_i, r_{t_i}^i(m))| - |P_{t_i}^i(m)| = (n-1)q^2$ for $i \in [q+1]$. This shows the proposition. $\square$

As $F(m)$ induces a clique and A_{ij}, B_h, R are subsets of $F(m)$, any two distinct vertices in any pair of these sets are adjacent.

Now we determine the number of neighbours of $w \in C_i$ where $i \in [q+1]$.

Lemma 5.7. *With the above notation, let $w \in C_i$ where $i \in [q+1]$. Then w has exactly*

- (i) $q(q-1)$ neighbours in C_j for $j \in [q+1] - \{i\}$,
- (ii) $q-1$ neighbours in A_{hj} if $i \in \{h, j\}$ and 0 otherwise,
- (iii) $q-1$ neighbours in B_j if $i = j$ and 0 otherwise,
- (iv) 0 neighbours in R .

Proof. Note that A_{ij}, B_i and C_i are subsets of $C(i, t_i, r_{t_i}^i(m))$ for $i, j \in [q+1]$ and $i \neq j$. This shows (iii) and (ii) for $i \in \{h, j\}$. Assume that $w = (\mathbf{y}, p)$. By Lemma 5.5, $N(w) \cap F(m)$ is the plane $P_{t_i}^i(m)$, and hence w has no neighbours in A_{jh} if $i \notin \{j, h\}$ by Theorem 2.6. This finishes the proof of (ii). It also follows that w has no neighbours in R , showing (iv). To show (i), let $j \in [q+1] \setminus \{i\}$. We have $r_{t_i}^i(p) = r_{t_i}^i(m)$, and thus $r_{t_j}^j(p) \neq r_{t_j}^j(m)$ by the definition of $TLS(q, n)$ and $\mathcal{O}$. We have $N(w) \cap F(m) = P_{t_i}^i(m)$, by Lemma 5.5, of which exactly q are in $C_{t_j}^j(m)$, namely the elements of $P_{t_i}^i(m) \cap P_{t_j}^j(m)$. As w has q^2 neighbours in $C(j, t_j, r_{t_j}^j(m))$ by Lemma 5.3, it follows that w has exactly $q^2 - q$ neighbours in C_j . This finishes the proof of the lemma. $\square$

Proposition 5.6 and Lemma 5.7 implies that there are four kinds of vertices in the local graph $\Delta(u)$. To make this precise:

Proposition 5.8. *With the above notation, let $a \in A_{ij}, b \in B_i, c \in C_i$, and $v \in R$ for $i, j \in [q+1]$. Then we have:*

- (i) $\lambda(u, a) = |F(m) - \{u, a\}| + |C_i| + |C_j| = q^3 - 2 + 2(n-1)q^2;$
- (ii) $\lambda(u, b) = |F(m) - \{u, b\}| + |C_i| = q^3 - 2 + (n-1)q^2;$
- (iii) $\lambda(u, c) = \sum_{t \neq i} |N(c) \cap C_t| + |C(i, t_i, r_{t_i}^i(m)) - \{u, c\}| = q(q^2 - q) + nq^2 - 2 = q^3 - 2 + (n-1)q^2;$

$$(iv) \lambda(u, v) = |F(m) - \{u, v\}| = q^3 - 2.$$

This gives immediately the following:

Proposition 5.9. *With the above notation, let $\mathcal{A} := \bigcup_i \bigcup_{j \neq i} A_{ij}$, $\mathcal{B} = \bigcup_i B_i$, and $\mathcal{C} := \bigcup_i C_i$. Then the partition $\{\mathcal{A}, \mathcal{B}, \mathcal{C}, R\}$ is an equitable partition of the local graph $\Delta(u)$ with quotient matrix*

$$Q := \begin{pmatrix} \binom{q+1}{2}(q-1) - 1 & (q+1)(q-1) & 2(n-1)q^2 & \frac{q(q-1)^2}{2} \\ \binom{q+1}{2}(q-1) & (q+1)(q-1) - 1 & (n-1)q^2 & \frac{q(q-1)^2}{2} \\ q(q-1) & q-1 & (n-1)q^2 + q(q^2 - q) - 1 & 0 \\ \binom{q+1}{2}(q-1) & (q+1)(q-1) & 0 & \frac{q(q-1)^2}{2} - 1 \end{pmatrix}.$$

As a consequence of the above two propositions we have the following result.

Proposition 5.10. *Each twisted Latin Square graph $TLS(q, n)$ is (α, β) -weakly-edge-regular, where $\alpha + 1 = (n-1)q^2 + q^3 - 2$ and $\beta = (2 - q^3) - q^2(n-1)(q^2 - q + 1)$.*

Proof. It can be directly confirmed by calculation, by using Propositions 5.8 and 5.9. $\square$

Proposition 5.11. *Each twisted Latin Square graph $TLS(q, n)$ is (μ, γ) -strongly co-edge-regular, where $\mu = (q+1)q^2$ and $\gamma = \mu(q^3 - 2 + (n-1)q^2)$.*

Proof. Let $u = (\mathbf{x}, m)$ be a vertex of the graph $TLS(q, n)$, where $\mathbf{x} \in \mathbb{F}_q^3$ and $m \in [n^2]$. We use the above notations A_{ij} , and so on. Let $w = (\mathbf{y}, m')$ be a non-adjacent vertex to u . Then $m' \neq m$. We have to consider two cases for w : either $N(w) \cap F(m) = \emptyset$ or $N(w) \cap F(m) \neq \emptyset$. In the first case, we have $N(w) \cap N(u) = N(w) \cap (\bigcup_{i=1}^{q+1} C_i)$. This implies $|N(w) \cap N(u)| = (q+1)q^2$ and

$$\sum_{v \in N(w) \cap N(u)} \lambda(u, v) = (q+1)q^2(q^3 - 2 + (n-1)q^2)$$

by Lemma 5.3 and Proposition 5.8.

If $N(w) \cap F(m) \neq \emptyset$, then, by Lemma 5.5, there exists $s \in [q+1]$ and $t \in [q]$ such that $N(w) \cap F(m) = P_t^s(m)$. We obtain that $|A_{ij} \cap P_t^s(m)| = 1$ for $i, j \in [q+1] - \{s\}$, and hence $|B_i \cap P_t^s(m)| = 1$ for $i \in [q+1] - \{s\}$. So that means $|\mathcal{A} \cap P_t^s(m)| = \frac{q(q-1)}{2}$ and $|\mathcal{B} \cap P_t^s(m)| = q$. We obtain that $|R \cap P_t^s(m)| = q^2 - \frac{q(q-1)}{2} - q = \frac{q(q-1)}{2}$. Also this implies that w has exactly $q^2 - q$ neighbours in C_i if $i \in [q+1] - \{s\}$ and exactly q^2 neighbours in C_s , by Lemma 5.3 and Proposition 5.8. As w has exactly q^2 neighbours in $F(m)$, we see that

$$|N(w) \cap N(u)| = |N(w) \cap F(m)| + \sum_{i=1}^{q+1} |N(w) \cap C_i| = q^2 + q^2 + q(q^2 - q) = (q+1)q^2.$$

Now we have

$$\begin{aligned} \sum_{v \in N(w) \cap N(u)} \lambda(u, v) &= \sum_{v \in N(w) \cap \mathcal{A}} \lambda(u, v) + \sum_{v \in N(w) \cap \mathcal{B}} \lambda(u, v) + \sum_{v \in N(w) \cap \mathcal{C}} \lambda(u, v) + \sum_{v \in N(w) \cap R} \lambda(u, v) \\ &= \frac{q(q-1)}{2}(q^3 - 2 + 2(n-1)q^2) + q(q^3 - 2 + (n-1)q^2) + \\ &\quad (q(q^2 - q) + q^2)(q^3 - 2 + (n-1)q^2) + \frac{q(q-1)}{2}(q^3 - 2) \\ &= (q+1)q^2(q^3 - 2 + (n-1)q^2). \end{aligned}$$

This shows the proposition. $\square$

Corollary 5.12. *Each twisted Latin Square graph $TLS(q, n)$ is a co-edge-regular graph of level 3 with spectrum*

$$\{(q^3 - 1 + q^2(q+1)(n-1))^{-1}, (q^2(n-1) - 1)^{(qn-1)(q+1)}, (-1)^{(q-1)q^2n^2}, (-q^2 - 1)^{(qn-1)(n-1)q}\}.$$

Proof. By Proposition 5.8 and Proposition 5.11, $TLS(q, n)$ is a co-edge-regular graph of level 3.

By Lemma 5.2 and Proposition 5.10, each twisted Latin Square graph $TLS(q, n)$ is k -regular where $k = q^3 - 1 + (q + 1)(n - 1)q^2$, and weakly-edge-regular with $\alpha = (n - 1)q^2 + q^3 - 3$ and $\beta = (2 - q^3) - q^2(n - 1)(q^2 - q + 1)$. By Proposition 5.11, $TLS(q, n)$ is strongly co-edge-regular with $\mu = (q + 1)q^2$ and $\gamma = \mu(q^2 - q)q + nq^2 - 2$. Note that $TLS(q, n)$ is not a strongly regular graph as there are three different valencies in any local graph of $TLS(q, n)$. Therefore, $TLS(q, n)$ has exactly four distinct eigenvalues $k = \theta_0 > \theta_1 > \theta_2 > \theta_3$ by Theorem 3.2.

It follows from Lemma 5.4 that $\theta_1 = q^2(n - 1) - 1$. By Theorem 3.2 and Equation (1), we obtain $\alpha - \mu = f_1 = \theta_1 + \theta_2 + \theta_3$ and $\mu(\alpha - 1) + k - \beta - \gamma = f_2 = \theta_1\theta_2 + \theta_1\theta_3 + \theta_2\theta_3$. Therefore, $\theta_2 = -1$ and $\theta_3 = -q^2 - 1$. The multiplicities can be determined by using $Tr(A) = 0$, $Tr(A^2) = q^3n^2k$, and that the sum of multiplicities is equal to $q^3n^2 = |V(TLS(q, n))|$. $\square$

As a consequence of the above corollary and Theorem 2.4, we have the following result which provides a proof of Theorem 1.5.

Theorem 5.13. *For a prime power q , there exist infinitely many integers n such that a twisted Latin Square graph $TLS(q, n)$ exists. The graph $TLS(q, n)$ is a co-edge-regular graph with level 3 and cospectral with the q -clique extension of each Latin Square graph $LS_{q+1}(qn)$.*

It worth mentioning that $TLS(q, n)$ is not switching equivalent to the q -clique extension of $LS_{q+1}(qn)$ if $n \geq 6$.

The smallest example of a twisted Latin Square graph is $TLS(2, 2)$, which has 32 vertices. Van Dam and Spence [19] provided tables of connected regular graphs with four distinct eigenvalues and at most 30 vertices.

In particular, we can also consider cliques-extensions of twisted Latin Square graphs. As a consequence, for any positive integer $s \geq 2$, there exist infinitely many co-edge-regular graphs that are non-isomorphic but cospectral with the s -clique extension of certain Latin Square graphs

Theorem 5.14. *Let q be a prime power. For any positive integer $s \geq 2$, such that q is a factor of s . If $q \neq s$, then there exist infinitely many integers n for which a co-edge-regular graph with level 4 exists that is cospectral with the s -clique extension of each Latin Square graph $LS_{q+1}(qn)$.*

Proof. By Theorem 5.13, there exists infinitely many integers n such that a twisted Latin Square graph $TLS(q, n)$ exists. Then the $\frac{s}{q}$ -clique extension of $TLS(q, n)$ is a co-edge-regular graph of level at least 3 with the same spectrum as the s -clique extension of $LS_{q+1}(qn)$. Moreover, if $s \neq q$, then the $\frac{s}{q}$ -clique extension of $TLS(q, n)$ is a co-edge-regular graph of level 4. $\square$

Remark 5.15. *Recently we were able to generalize the construction in this paper to obtain an infinite family of co-edge-regular graphs with unbounded level that are cospectral with clique-extensions of Latin square graphs. A detailed exposition of these results will be given in a forthcoming work.*

We conclude with some open problems:

- Problem 5.16.** (i) *Let G be a connected co-edge-regular graph with level t and with four distinct eigenvalues and smallest eigenvalue $-\theta$. Is t bounded by a function of θ ?*
- (ii) *Is it possible to construct infinite families of cospectral graphs of certain clique extensions of certain Steiner graphs in a similar manner as in Section 5?*
- (iii) *Is it possible to generalize Theorem 1.1 to the class of co-edge-regular graphs of level 2 with exactly four distinct eigenvalues?*

Acknowledgements

J.H. Koolen is partially supported by the National Key R. and D. Program of China (No. 2020YFA0713100), the National Natural Science Foundation of China (No. 12071454, No. 12471335), and the Anhui Initiative in Quantum Information Technologies (No. AHY150000). We thank Dr. Qianqian Yang for her helpful comments.

References

- [1] E. Bannai, E. Bannai, T. Ito, and R. Tanaka. *Algebraic Combinatorics*. De Gruyter, Berlin, Boston, 2021.
- [2] R. C. Bose. An affine analogue of Singer’s theorem. *J. Indian Math. Soc. (N.S.)*, 6:1–15, 1942.
- [3] A. E. Brouwer and W. H. Haemers. *Spectra of Graphs*. Springer, New York, 2012.
- [4] A. E. Brouwer and H. Van Maldeghem. *Strongly regular graphs*. Cambridge University Press, Cambridge, 2022.
- [5] C. Godsil and G. F. Royle. *Algebraic graph theory*, volume 207. Springer Science & Business Media, 2001.
- [6] F. Goldberg. On quasi-strongly regular graphs. *Linear Multilinear Algebra*, 54(6):437–451, 2006.
- [7] W. H. Haemers. Hoffman’s ratio bound. *Linear Algebra and its Applications*, 617:215–219, 2021.
- [8] W. H. Haemers and V. D. Tonchev. Spreads in strongly regular graphs. *Des. Codes Cryptogr.*, 8(1-2):145–157, 1996.
- [9] S. Hayat, J. H. Koolen, and M. Riaz. A spectral characterization of the s -clique extension of the square grid graphs. *European Journal of Combinatorics*, 76:104–116, 2019.
- [10] A.J. Hoffman. On the polynomial of a graph. *Amer. Math. Monthly*, 70:30–36, 1963.
- [11] A. Jurišić, J. H. Koolen, and P. Terwilliger. Tight distance-regular graphs. *J. Algebraic Combin.*, 12(2):163–197, 2000.
- [12] H. F. MacNeish. Euler’s squares. *Ann. of Math.*, 23:221–227, 1922.
- [13] A. Neumaier. Strongly regular graphs with smallest eigenvalue $-m$. *Arch. Math. (Basel)*, 33(4):392–400, 1979/80.
- [14] D. K. Ray-Chaudhuri and R. M. Wilson. The existence of resolvable block designs. In *A survey of combinatorial theory (Proc. Internat. Sympos., Colorado State Univ., Fort Collins, Colo., 1971)*, pages 361–375. North-Holland, Amsterdam-London, 1973.
- [15] Y. Tan, J. H. Koolen, and Z. Xia. A spectral characterization of the s -clique extension of the triangular graphs. *Discuss. Math. Graph Theory*, 40(2):663–676, 2020.
- [16] J. A. Thas and S. E. Payne. Spreads and ovoids in finite generalized quadrangles. *Geom. Dedicata*, 52(3):227–253, 1994.
- [17] E. R. van Dam. Regular graphs with four eigenvalues. *Linear Algebra Appl.*, 226–228:139–162, 1995.
- [18] E. R. van Dam. Three-class association schemes. *J. Algebraic Combin.*, 10(1):69–107, 1999.
- [19] E. R. van Dam and E. Spence. Small regular graphs with four eigenvalues. *Discrete Math.*, 189(1-3):233–257, 1998.